


Measures to limit the spread of the flu:


Severe Pandemic:

All of the seasonal and mild to moderate flu methods plus:

- Possible extended school closure, which could range from weeks to up to 3 months. Presently, the Department of Health and Human Services, the Centers for Disease Control and Prevention, the Department of Education, and other agencies are conducting a comprehensive review of school closure and its feasibility and effects in mitigating a severe pandemic. The results of this study will be shared in the coming months.
- Promote social distancing of children and teens outside the school setting by reducing their social circulation and contacts to the greatest extent possible. This could include canceling extracurricular activities.



School Closure



Social Distancing

Mild to Moderate Pandemic:

All of the seasonal flu methods plus the following:

- Encourage the use of social distancing at the work place, at school and in the community.
- Possible school closure for a short amount of time (possibly days to a couple of weeks).
- Work with community flu-planning team to assess whether any additional measures should be taken.



Assess Measures

Seasonal Flu:

- Promote hand washing and cough hygiene via school-wide campaigns and modeling by school staff.
- Encourage vaccination of staff and students for whom the flu vaccine is recommended.
- Persons developing symptoms at school should be sent home as soon as possible and instructed by appropriate officials not to return until they are well.



Promote hygiene



Encourage vaccination



Stay at home

Infection control

The primary strategies for preventing spread of pandemic influenza:

Persons with flu symptoms should:

- Stay at home;
- Cover nose and mouth when coughing or sneezing;
- Wash hands with soap and water or use alcohol-based hand sanitizers frequently; and
- Try to maintain spatial separation of at least three feet from others if possible.

Hand, Cough and Sneeze Hygiene

- When sneezing or coughing, cover the nose and mouth with a tissue or upper arm if a tissue is not available.
- Dispose of used tissues in a wastebasket and wash hands after coughing, sneezing, or blowing nose.
- Use warm water and soap or alcohol-based hand sanitizers to clean hands.
- Wash hands before eating or touching eyes, nose, or mouth.



Social distancing

In a pandemic, the risk of getting the flu is greatest when one has close contact with an infected person. Social distancing measures such as school closure, telecommuting or staggered shifts for the workforce, and cancellation of public gatherings may be effective in reducing transmission risks.

